arXiv:1212.6404v1 [hep-ex] 27 Dec 2012

Search for $D^0 \to \ell^+\ell^-$ and $D^0 \to \gamma\gamma$ at *BaBar* and BESIII

Alberto Lusiani (on behalf of the BaBar and BESIII collaborations)
Scuola Normale Superiore e INFN di Pisa

Proceedings of CKM 2012, the 7th International Workshop on the CKM Unitarity Triangle, University of Cincinnati, USA, 28 September - 2 October 2012

1 Introduction

Flavour changing neutral currents (FCNC) leading to the decay of charmed D mesons into $\ell^+\ell^-$ and $\gamma\gamma$ are very suppressed in the Standard Model (SM), and in fact the Glashow-Iliopoulos-Maiani (GIM) mechanism [1] is more effective here than for B and K mesons because the down-type quark mass differences are relatively smaller than the up-type ones. In the SM, the $D^0 \to \mu^+\mu^-$ and $D^0 \to e^+e^-$ branching fractions are dominated by long-distance contributions and estimated to be $\mathcal{B}(D^0 \to \mu^+\mu^-) \sim 10^{-13}$ and $\mathcal{B}(D^0 \to e^+e^-) \sim 10^{-23}$ [2], while $D^0 \to e^\pm\mu^\mp$ violates lepton flavour conservation and is forbidden in the SM. If neutrino mixing is accomodated in the SM, $D^0 \to e^\pm\mu^\mp$ would anyway happen only at undetectable small rates. The SM predictions are all well below the present experimental sensitivities, but searches for these decays can probe NP contributions that may provide enhancements of several orders of magnitude [3]. Furthermore, NP models can affect $D^0 \to \ell^+\ell^-$ and $D^0 - \overline{D^0}$ mixing in a correlated way [3]. The short-distance SM contribution to $D^0 \to \gamma\gamma$ is $\simeq 3{\cdot}10^{-11}$, the dominant long-distance contribution is $\simeq 3.5{\cdot}10^{-8}$ [4]. NP models can enhance the SM rate by more than 2 orders of magnitude [5].

2 *BaBar* search for $D^0 \to \ell^+\ell^-$

BaBar has searched for $D^0 \to \ell^+\ell^-$ [6] using approximately 468 fb^{-1} of e^+e^- collisions produced by the PEP-II asymmetric-energy e^+e^- collider at and near the $\Upsilon(4S)$ resonance.

D^0 candidates are found by combining pairs of oppositely charged leptons. Decays to $\pi^+\pi^-$ and $K^-\pi^+$ are used as control samples. D mesons from direct $e^+e^- \to c\overline{c}$ production are selected by requiring that the D^0 candidate momentum be larger than 2.4 GeV/c, which is about the kinematic limit for $B \to D^*\pi$, $D^{*+} \to D^0\pi^+$. Combinatorial background is suppressed by requiring that the D^0 candidate originate from the decay $D^*(2010)^+ \to D^0\pi^+$ [1]. A fit of the $D^{*+} \to D^0\pi^+; D^0 \to \ell^+\ell^-$ decay chain is performed where the D^0 tracks are constrained to come from a common vertex and the D^0 and slow pion are constrained to form a common vertex within the beam interaction region. The χ^2 probabilities of the D^0 and D^* vertices from this fit must be at least 1%. The reconstructed D^0 mass $m(D^0)$ must be within $[1.65, 2.05]$ GeV/c^2 and the mass difference Δm must be within $[0.141, 0.149]$ GeV/c^2. The D^0 and Δm mass resolutions, measured in the $D^0 \to \pi^+\pi^-$ sample, are 8.1 MeV/c^2 and 0.2 MeV/c^2, respectively.

Background candidates are either random combinations of two leptons (combinatorial background), or $D^0 \to \pi^+\pi^-$ decays where both pions pass the lepton identification criteria (peaking background). The $D^0 \to \pi^+\pi^-$ background is most important for the $D^0 \to \mu^+\mu^-$ channel. Combinatorial background from semileptonic B decays is suppressed by using a Fisher discriminant [7] of five variables. Events with hard initial state radiation that converts to eletron pairs are suppressed by requiring at least 5 tracks for the $D^0 \to e^+e^-$ channel and at least 4 tracks for the $D^0 \to \mu^+\mu^-$ and $D^0 \to e^\pm\mu^\mp$ channels, by accepting at most 3 electron candidates and by vetoing particles compatible with being results of photon conversions. The selection criteria for each signal channel were chosen to give the lowest expected signal branching fraction upper limit for the null hypothesis (a true branching fraction of zero) using the MC samples.

Control samples and sidebands are used to estimate backgrounds. Combinatorial background is estimated using the upper sideband in the D^0 mass distribution. Peaking background is estimated by scaling the observed $D^0 \to \pi^+\pi^-$ yield with the misidentification rates estimated from the $D^0 \to K^-\pi^+$ data control sample.

For all three channels, no statistically significant yield is observed compared with the expected background: 1 event observed vs. 1.0 ± 0.5 expected background for $D^0 \to e^+e^-$, 2 observed vs. 1.4 ± 0.3 expected for $D^0 \to e^\pm\mu^\mp$ and 8 observed vs. 3.9 ± 0.6 expected for $D^0 \to \mu^+\mu^-$. The probability of observing 8 events when 3.9 ± 0.6 events are expected is 5.4%. Accounting also for systematic uncertainties, the 90% confidence level (CL) upper limits for the branching fractions are $\mathcal{B}(D^0 \to e^+e^-) < 1.7 \cdot 10^{-7}$, $\mathcal{B}(D^0 \to e^\pm\mu^\mp) < 3.3 \cdot 10^{-7}$ and $\mathcal{B}(D^0 \to \mu^+\mu^-) < 2.6 \cdot 10^{-7}$ [6].

3 *BABAR* search for $D^0 \to \gamma\gamma$

The *BABAR* collaboration has conducted a search for $D^0 \to \gamma\gamma$ [8] on 470.5 fb^{-1} of data collected by the *BABAR* detector at the SLAC PEP-II e^+e^- asymmetric-energy collider operating at e^+e^- CM energies at and around the $\Upsilon(4S)$ peak. The important background decay mode $D^0 \to \pi^0\pi^0$ and the decay mode $D^0 \to K^0_S\pi^0$ are also selected and studied in the same analysis. The last channel is used as reference normalization mode to avoid uncertainties in the number of D^{*+}.

---

[1] Here and in the following charge conjugation is implied unless otherwise stated

D^0 candidates are made with pairs of photons and pairs of π^0 mesons, and the resulting D^0 is required to come from the decay $D^*(2010)^+ \to D^0\pi^+$ (D^{*+} tag). The candidates are required to have a momentum $> 2.85\,\mathrm{GeV}/c$ and their invariant mass must be between 1.7 and $2.1\,\mathrm{GeV}/c^2$ for the $D^0 \to \gamma\gamma$ analysis and between 1.65 and $2.05\,\mathrm{GeV}/c^2$ for the $D^0 \to \pi^0\pi^0$ analysis. The photon candidates are selected to have CM energies between 0.74 and 4 GeV, the π^0 candidates must have energy larger than 0.6 GeV. The D^0 candidates for the $D^0 \to K^0_S\pi^0$ reference mode are formed by combining a π^0 candidate as defined above with a K^0_S candidate consistent with the decay $K^0_S \to \pi^+\pi^-$, with a decay length significance (flight length divided by its estimated uncertainty) larger than 3. The $D^0 \to \gamma\gamma$ selection rejects events whose photons are compatible with a π^0 decay. A kinematic fit is applied to the events, requiring the candidate D^0 invariant mass to be between 1.6 and $2.1\,\mathrm{GeV}/c^2$. Both the D^0 and π^+ are constrained to originate from a common vertex within the beamspot to satisfy the D^{*+} tag requirement. The $D^0 \to \gamma\gamma$ analysis signal efficiency is 6.1% with the corresponding reference mode ($D^0 \to K^0_S\pi^0, K^0_S \to \pi^+\pi^-$) efficiency at 7.6%. The $D^0 \to \pi^0\pi^0$ analysis signal and reference mode efficiencies are 15.2% and 12.0%, respectively.

For all modes, signal yields are determined using unbinned maximum likelihood fits to the invariant mass distribution of D^0 candidates. The probability distribution functions (PDFs) for signal and backgrounds have their shapes fixed from Monte Carlo simulation and their relative normalization fit on data. The invariant $\gamma\gamma$ mass distribution of the $D^0 \to \gamma\gamma$ candidates is shown in Fig. 1 together with the fit of the signal and background component contributions. The signal yield is -6 ± 15, consistent with no $D^0 \to \gamma\gamma$ events. Using the $D^0 \to K^0_S\pi^0$ yield, $\mathcal{B}(D^0 \to \gamma\gamma) = (-0.49 \pm 1.23 \pm 0.02)\cdot 10^{-6}$ where the errors are the statistical uncertainty and the uncertainty in the reference mode branching fraction, respectively.

Several systematic uncertainties cancel partially or completely by normalizing the yield to the reference mode; the largest estimated systematic contributions come from uncertainties in tracking efficiency (1.4%), π^0 veto (1.8%), photon reconstruction (0.6%). The π^0 veto performance has been studied on data using the physically forbidden decay $D^0 \to K^0_S\gamma$. Accounting for systematic uncertainties, the 90% CL upper limit for B ($D^0 \to \gamma\gamma$) is determined to be $< 2.4 \cdot 10^{-6}$ [8].

4 BESIII search for $D^0 \to \gamma\gamma$

The BESIII collaboration has reported preliminary results on a search for $D^0 \to \gamma\gamma$ [9] based on an integrated luminosity of $\sim$2.9 fb^{-1} of e^+e^- collisions produced at the BEPCII energy-symmetric ring at $\sqrt{s} = 3.773$ GeV and recorded with the BESIII



Figure 1: The $\gamma\gamma$ mass distribution for $D^0 \to \gamma\gamma$ candidates in data (data points). The solid blue curve corresponds to the signal plus background fit, where the signal component has a slight negative yield, the long-dash red curve corresponds to combinatorial background component, and the small-dash pink curve corresponds to the combinatoric background plus $D^0 \to \pi^0\pi^0$ background shape. The χ^2 value is determined from binned data. The pull distribution shows differences between the data and the solid blue curve with values and errors normalized.

detector [10]. The analysis includes also the measurement of the major background $D^0 \to \pi^0\pi^0$.

To select the $D^0 \to \pi^0\pi^0$ channel, photons are first paired to form π^0 candidates, which are then re-fitted with a mass constraint. π^0 candidates are combined to form D^0 candidates, whose energy is required to match half the event energy with a discrepancy -60 MeV $< \Delta E <$ 30 MeV, where $\Delta E = E_{\pi^0\pi^0} - E_{\text{beam}}$. The signal yield is obtained with a maximum-likelihood fit to the beam-constrained mass $M_{bc} \equiv \sqrt{E^2_{\text{beam}} - p^2_D c^2}$ where E_{beam} is the beam energy and p_D is the $D^0(\to \pi^0\pi^0)$ candidate momentum. The fit yields 4081 ± 117 signal events.

The selection of $D^0 \to \gamma\gamma$ candidates starts with pairing the two most energetic photons. Events with photons compatible with $\pi^0 \to \gamma\gamma$ are rejected. Radiative Bhabha events are suppressed by requiring that photons be isolated at least by 20° from all tracks and by requiring that the most energetic track fail the electron identification.

Finally, the signal yield is extracted with a fit to the ΔE distribution after requiring that $1860 < M_{bc} < 1870$ MeV $/c^2$. Here, the signal shape is fixed from Monte Carlo. The fit has χ^2 of 63.7 for 76 degrees of freedom and determines a signal yield of -2.9 ± 7.1.

No significant signal is found and 90% CL upper limits are set to $\mathcal{B}(D^0 \to \gamma\gamma)/\mathcal{B}(D^0 \to \pi^0\pi^0) < 5.8{\cdot}10^{-3}$ and $\mathcal{B}(D^0 \to \gamma\gamma) < 4.7{\cdot}10^{-6}$ (preliminary) [9].

5 Conclusions

There is no evidence of New Physics in the searches that *BABAR* and BESIII have done for the rare charm decays $D^0 \to \ell^+\ell^-$ and $D^0 \to \gamma\gamma$. The experimental sensitivities are still far from the SM predictions, leaving space for further significant improvements.

References

[1] S.L. Glashow, J. Iliopoulos and L. Maiani, Phys. Rev. D2 (1970) 1285.

[2] G. Burdman et al., Phys.Rev. D66 (2002) 014009, arXiv:hep-ph/0112235 [hep-ph].

[3] E. Golowich et al., Phys.Rev. D79 (2009) 114030, arXiv:0903.2830 [hep-ph].

[4] G. Burdman and I. Shipsey, Ann.Rev.Nucl.Part.Sci. 53 (2003) 431, arXiv:hep-ph/0310076 [hep-ph].

[5] S. Prelovsek and D. Wyler, Phys.Lett. B500 (2001) 304, arXiv:hep-ph/0012116 [hep-ph].

[6] BaBar, J. Lees et al., Phys.Rev. D86 (2012) 032001, arXiv:1206.5419 [hep-ex].

[7] R.A. Fisher, Annals of Eugenics 7 (1936) 179.

[8] BaBar, J. Lees et al., Phys.Rev. D85 (2012) 091107, arXiv:1110.6480 [hep-ex].

[9] H. Muramatsu, arXiv:1208.4744 [hep-ex].

[10] M. Ablikim et al., Nuclear Instruments and Methods in Physics Research A 614 (2010) 345, arXiv:0911.4960 [physics.ins-det].